




Hurricanes and Health Providers

Hurricanes and Climate Change

Hurricanes – intense low-pressure tropical storms producing destructive wind speeds and heavy rain – have far-reaching effects on health, especially for individuals with chronic medical conditions. The likelihood of hurricanes making landfall in the United States is illustrated in the map below.



Map of hurricane risk in the United States. Courtesy of FEMA

Climate change warms the oceans, and hurricane strength comes from ocean heat. Climate models project more intense hurricanes and associated coastal flooding and wind damage in this century.¹ Trends show that hurricane peak intensity has been occurring closer to shore and their tracks have been moving north, putting more northern coastal areas, which may have less investment in preparedness, at greater risk from hurricane strikes.^{2,3}

Health Impacts of Hurricanes

Hurricanes affect health through strong winds, heavy rains, and storm surge, all of which can cause injuries. They also can result in floods that can disrupt essential services, including power, water, and healthcare access.

Hurricanes can increase morbidity and mortality for individuals with a variety of medical conditions, as well as affect perinatal outcomes.

- **Respiratory disease** related mortality has been found to increase 8.3% one month after hurricanes.⁴
- **Cardiovascular disease** hospitalizations were double in older patients up to 1 year after Hurricane Sandy. Myocardial infarctions and strokes were 22%⁵ and 7% higher, respectively, in areas more severely affected by the storm.⁶
- **Type II diabetes** related emergency room visits increased 84% during the week of Hurricane Sandy.⁷ Increased diabetic morbidity after hurricanes may result from lack of access to medications and healthy foods.⁸

- **Chronic kidney disease** patients on dialysis have elevated mortality after hurricanes and may be more likely to need emergency department care, be hospitalized, and receive dialysis during hospitalization.^{9,10} Risks to renal patients may result from lack of access to dialysis, concurrent heat wave exposure and associated health risks, infectious diseases, or direct trauma from the storm.⁹
- **Eye injuries** and infections may be more likely after hurricanes.¹¹
- **Pregnant** people exposed to hurricanes were more likely to experience **preterm birth**.¹² Stress associated with many forms of disaster has been linked to preterm birth, and this may be mediated through corticotropin releasing hormone and the HPA axis.¹³

Populations at greater risk for harm after hurricanes

Population	Risk
Black and Hispanic	<p>More likely to live:</p> <ul style="list-style-type: none"> • near industrial facilities that release toxic substances during and after hurricanes¹⁴ • in areas with high flood risk¹⁵
Chronic Diseases	<p>May be more likely to have disease flares due to:</p> <ul style="list-style-type: none"> • supply chain disruptions and medication shortages • difficulty accessing treatment or care • power outages affecting electric medical devices¹⁶
Disabilities	<p>Face barriers to:</p> <ul style="list-style-type: none"> • receiving emergency communications • evacuation due to inaccessible transportation options or lack of necessary mobility assistance • access to evacuation shelters¹⁷ <p>May be more likely to live in public housing or in higher flood risk areas.¹⁸</p>
Low-income	<p>More likely to live:</p> <ul style="list-style-type: none"> • closer to industrial facilities and toxic waste sites that may release pollution after a storm.¹⁹ • in areas with high flood risk^{14,15}
Older Adults	<p>More likely to have:</p> <ul style="list-style-type: none"> • medical comorbidities • limited capacity to evacuate or be resilient to a hurricane or flood, especially if they lose their homes^{20,21}
Rescue Workers and First Responders	<p>More likely to:</p> <ul style="list-style-type: none"> • sustain unintentional injuries during and after hurricanes¹⁶

Immediate health risks from hurricanes

Immediate impacts of hurricanes are often due to high windspeeds and flooding. Unintentional injuries are also common during and immediately following hurricanes, as well as during rebuilding efforts.

Common unintentional injuries due to hurricanes include:

- Drowning
- Electrocutation
- Cuts, lacerations, and puncture wounds
- Falls
- Traumatic head injuries from falling debris
- Trauma from motor vehicle accidents²²

Deaths from injuries after hurricanes have been found to peak about 1 month after a storm occurs.⁴

Infectious diseases associated with flooding

Flooding has been found to increase risks of infections, including vector, rodent and waterborne diseases, as well as illnesses associated with exposure to fungi. Infectious disease risk after hurricanes appears to peak 2 months after the event, but infections can be seen in the days after a storm occurs.⁴

Timing of post-flood infectious diseases in the United States

Early (<10 d after event)

- Cellulitis, including from *Vibrios*
- Pneumonias (may be aspiration related and polymicrobial)
- Viral respiratory infections
- Gastroenteritis (SSYCE, *Vibrios*, viruses)

Late (>10 d after event)

- Mosquito-borne illnesses
- Skin infection from atypical organisms (fungi, mycobacteria, mold)
- Hepatitis A or E

Adapted from Paterson et al. Clin Inf Dis. doi:10.1093/cid/ciy227 and Ivers and Ryan. Curr Op Inf Dis. doi: 10.1097/01.qco.0000244044.85393.9e.

Floods have been commonly associated with outbreaks of waterborne diseases, including from bacterial and viral pathogens.²³ Individuals who obtain water from private wells may be at particularly high risk, but even public water systems can be contaminated during extreme storms.²⁴

Standing water may create breeding grounds for disease-carrying mosquitos²⁵ and fungal growth. Flooded homes are much more likely to have higher levels of molds, such as *Aspergillus*, *Penicillium*, and *Cladosporium*.²⁶ Patients with allergic disorders may be more likely to develop symptoms from mold exposures.²⁷ Fungal infections are less common, but immunocompromised individuals have developed respiratory fungal infections after flood induced mold exposures.²⁷

Population displacement that occurs after hurricanes and floods can result in overcrowded homes and shelters, sometimes with inadequate sanitation. These conditions can result in the spread of many different infections, especially those transmitted by respiratory or fecal-oral routes.²⁷

Toxic exposures

Hurricanes have caused massive releases of toxic industrial, wastewater, and agricultural substances into the environment. What toxins may be most prevalent in a community after a flood may reflect what is stored in local toxic substance impoundments. The EPA tracks sites in your community that may contain hazardous substances which can be mobilized by flooding [here](#).ⁱ

Hurricanes Katrina and Harvey damaged fossil fuel processing and storage facilities, which released hydrocarbons (known to cause respiratory irritation and arrhythmias), volatile organic compounds (which can be carcinogenic), and heavy metals, such as lead, into air, water, and soil. Hurricane Florence breached a coal ash pit in North Carolina. Coal ash is a byproduct of coal combustion and coal ash pits are present across the country. Coal fly ash contains volatile organic compounds as well as heavy metals, including mercury, arsenic and/or chromium.

The figure below shows the location of coal ash contamination sites as of February 2014. Red denotes a coal ash contamination site. Green denotes a coal ash spill. Black denotes both a contaminated site and spill.



Sources: Esri, USGS | Esri, HERE, Garmin, FAO, NOAA, USGS, EPA-

Infrastructure failures can cause additional unique and harmful exposures. Gasoline shortages after Hurricane Sandy led to increased gastrointestinal and pulmonary symptoms due to gasoline-siphoning related exposures.²⁸ Acute intoxication from gasoline fumes can cause confusion, giddiness, nausea, headache, blurred vision, dizziness, and weakness. Severe exposure can result in respiratory depression, seizures, loss of consciousness, and coma.

Carbon monoxide poisonings often increase after disasters, most often from indoor or inadequately ventilated electrical generator use combined with absent or non-functional carbon monoxide monitors.²⁹ Patients may also attempt to cook or heat their homes by burning fuels, such as wood or propane, indoors, or to run their vehicle for air conditioning in an enclosed garage, which can also lead to carbon monoxide poisoning.

i. <https://cimc.epa.gov/ords/cimc/f?p=cimc:map::::71>

Hurricanes and mental health

Increases in PTSD, depression, and anxiety symptoms have been documented after hurricanes.¹⁶ Prevalence of diagnoses after hurricanes vary with the intensity of the storm. A meta-analysis of tropical cyclone survivors globally found ~18% had PTSD with more severe storms.³⁰ Need for psychiatric care can extend for prolonged periods after hurricanes make landfall. Reports of psychiatric concerns have been elevated for years following Hurricanes Sandy and Katrina.^{31–33}

Children, females, and older individuals may be most at risk of mental health symptoms after hurricanes.³⁴ Individuals in certain occupations, such as farm workers and first responders (e.g., emergency medical personnel, police officers, and fire fighters), may also be more at risk. Mental health symptoms after exposure to a hurricane have also been found to be associated with lack of necessities, including medications and access to medical care as discussed above, personal physical exposure to hurricane forces (e.g., wind, rain, storm surge), evacuation, loss of a loved one or pet, repeat exposure to hurricanes, existing mental health disorders or recent personal adversity.^{35–37}

Disruption of health-systems infrastructure and displacement

Hurricanes often impede healthcare delivery as they can damage healthcare facilities, cut off essential utilities, disrupt supply chains, and inundate roads. Patients who are displaced by storms may be unable to refill medications, and lack of access to medications after hurricanes and floods has been associated with increased morbidity.^{22,38}

Patients with chronic conditions, including diabetes and cancer, may delay necessary healthcare services because of healthcare facility closure, difficulty with transportation, or competing demands on time²⁰ and may have worse outcomes as a result.³⁹

Hurricane Action Plans for Patients

We recommend that you familiarize yourself with the “Hurricane Action Plan” provided in the toolkit and review it with any patient at risk of experiencing a hurricane. The action plan can be provided during care visits with adolescents and adults and can be the basis for a discussion around safety planning and care management in the event of a hurricane. Action plans should be completed before hurricane season in your locale.

Anticipatory Guidance for Providers to Give to Patients

Anticipatory guidance for hurricanes may contribute to improved health outcomes. The strategies and resources below may be helpful for you to provide to patients who are at risk from hurricanes and reflect the “Hurricane Action Plan” available in the toolkit.

Forecasts

Baseline and future flooding risk for many properties in the United States can be found at riskfactor.com. Hurricane forecasts are available from several outlets including the [National Hurricane Center](http://www.nhc.noaa.gov/).ⁱⁱ

A “hurricane watch” indicates that hurricane conditions (winds of greater than 74 mph) are possible. A “hurricane warning” indicates that hurricane conditions are expected. Hurricane warnings are given 36 hours prior to the expected hurricane impact to give residents time to prepare or evacuate.

ii. <https://www.nhc.noaa.gov/>

References

1. Knutson TR, Sirutis JJ, Bender MA, Tuleya RE, Schenkel BA. Dynamical downscaling projections of late twenty-first-century U.S. landfalling hurricane activity. *Climatic Change* 2022 171:3. 2022;171(3):1-23. doi:10.1007/S10584-022-03346-7
2. Baldini LM, Baldini JUL, McElwaine JN, et al. Persistent northward North Atlantic tropical cyclone track migration over the past five centuries. *Sci Rep*. 2016;6(1):37522. doi:10.1038/srep37522
3. Wang S, Toumi R. Recent migration of tropical cyclones toward coasts. *Science*. 2021;371(6528):514-517. doi:10.1126/SCIENCE.ABB9038
4. Parks RM, Benavides J, Anderson GB, et al. Association of Tropical Cyclones With County-Level Mortality in the US. *JAMA*. 2022;327(10):946-955. doi:10.1001/JAMA.2022.1682
5. Dresser C, Balsari S, Leaning J. Hurricanes and Health. *Oxford Research Encyclopedia of Natural Hazard Science*. Published online March 23, 2022. doi:10.1093/ACREFORE/9780199389407.013.359
6. Swerdel JN, Janevic TM, Cosgrove NM, Kostis JB. The effect of Hurricane Sandy on cardiovascular events in New Jersey. *J Am Heart Assoc*. 2014;3(6). doi:10.1161/JAHA.114.001354
7. Velez-Valle E, Shendell D, Health SE... of E, 2016 undefined. Type II diabetes emergency room visits associated with Hurricane Sandy in New Jersey: implications for preparedness. *JSTOR*. 2016;79(2):30-37. Accessed September 2, 2022. <https://www.jstor.org/stable/26330515>
8. Travia KR, Kahkoska AR, Igudesman D, Souris KJ, Beasley C, Mayer-Davis EJ. Impact of Hurricane Matthew on Diabetes Self-Management and Outcomes. *N C Med J*. 2021;82(2):100. doi:10.18043/NCM.82.2.100
9. Blum MF, Feng Y, Anderson GB, Segev DL, McAdams-DeMarco M, Grams ME. Hurricanes and Mortality among Patients Receiving Dialysis. *Journal of the American Society of Nephrology*. 2022;33(9):1757-1766. doi:10.1681/ASN.2021111520
10. Kelman J, Finne K, Bogdanov A, et al. Dialysis Care and Death Following Hurricane Sandy. *American Journal of Kidney Diseases*. 2015;65(1):109-115. doi:10.1053/J.AJKD.2014.07.005
11. Karimaghaei C, Merkley K, Nazari H. Ophthalmology emergency room admission after Hurricane Harvey. *Am J Disaster Med*. 2021;16(4):255-261. doi:10.5055/AJDM.2021.0409
12. Sun S, Weinberger KR, Yan M, Brooke Anderson G, Wellenius GA. Tropical cyclones and risk of preterm birth: A retrospective analysis of 20 million births across 378 US counties. *Environ Int*. 2020;140:105825. doi:10.1016/J.ENVINT.2020.105825
13. Ng PC. The fetal and neonatal hypothalamic–pituitary–adrenal axis. *Arch Dis Child Fetal Neonatal Ed*. 2000;82(3):F250-F254. doi:10.1136/FN.82.3.F250
14. Perlin SA, Wong D, Sexton K. Residential proximity to industrial sources of air pollution: interrelationships among race, poverty, and age. *J Air Waste Manag Assoc*. 2001;51(3):406-421. doi:10.1080/10473289.2001.10464271
15. Bakkensen LA, Ma L, Blomquist G, et al. Sorting Over Flood Risk and Implications for Policy Reform. Published online July 2020. Accessed September 3, 2022. https://www.frbsf.org/economic-research/wp-content/uploads/sites/4/Bakkensen_Ma_2020.pdf
16. Waddell SL, Jayaweera DT, Mirsaeidi M, Beier JC, Kumar N. Perspectives on the Health Effects of Hurricanes: A Review and Challenges. Published online 2021. doi:10.3390/ijerph18052756
17. Powell R, Gilbert S. *The Impact of Hurricanes Katrina and Rita on People with Disabilities: A Look Back and Remaining Challenges*.; 2006. Accessed September 3, 2022. <https://ncd.gov/publications/2006/aug072006>
18. Chakraborty J, McAfee AA, Collins TW, Grineski SE. Exposure to Hurricane Harvey flooding for subsidized housing residents of Harris County, Texas. *Natural Hazards*. 2021;106(3):2185-2205. doi:10.1007/S11069-021-04536-9/TABLES/4
19. Friedrich MJ. Determining Health Effects of Hazardous Materials Released During Hurricane Harvey. *JAMA*. 2017;318(23):2283-2285. doi:10.1001/JAMA.2017.15558
20. Wang C, Bowers BJ. Bracing For Hurricanes: A Qualitative Analysis of the Extent and Level of Preparedness Among Older Adults. *Gerontologist*. 2018;58(1):57-67. doi:10.1093/GERONT/GNX187
21. Prohaska TR, Peters KE. Impact of Natural Disasters on Health Outcomes and Cancer Among Older Adults. *Gerontologist*. 2019;59(Supplement_1):S50-S56. doi:10.1093/GERONT/GNZ018

22. Lee S, Jayaweera DT, Mirsaeidi M, Beier JC, Kumar N. Perspectives on the Health Effects of Hurricanes: A Review and Challenges. *Int J Environ Res Public Health*. 2021;18(5):1-18. doi:10.3390/IJERPH18052756
23. Levy K, Woster AP, Goldstein RS, Carlton EJ. Untangling the Impacts of Climate Change on Waterborne Diseases: A Systematic Review of Relationships between Diarrheal Diseases and Temperature, Rainfall, Flooding, and Drought. *Environ Sci Technol*. 2016;50(10):4905-4922. doi:10.1021/ACS.EST.5B06186/ASSET/IMAGES/LARGE/ES-2015-06186D_0005.JPEG
24. Mapili K, Rhoads WJ, Coughter M, Pieper KJ, Edwards MA, Pruden A. Occurrence of opportunistic pathogens in private wells after major flooding events: A four state molecular survey. *Science of The Total Environment*. 2022;826:153901. doi:10.1016/J.SCITOTENV.2022.153901
25. Centers for Disease Control and Prevention. Mosquitoes, Hurricanes, and Flooding. Accessed August 31, 2022. <https://www.cdc.gov/mosquitoes/mosquito-control/community/mosquitoes-and-hurricanes.html>
26. Barbeau DN, Grimsley LF, White LE, El-Dahr JM, Lichtveld M. Mold Exposure and Health Effects Following Hurricanes Katrina and Rita. *Annu Rev Public Health*. 2010;31:165-178. doi:10.1146/annurev.publhealth.012809.103643
27. Saulnier DD, Brolin Ribacke K, von Schreeb J. No calm after the storm: a systematic review of human health following flood and storm disasters. *Prehosp Disaster Med*. 2017;32(5):1-12. doi:10.1017/s1049023x17006574
28. Kim HK, Takematsu M, Biary R, Williams N, Hoffman RS, Smith SW. Epidemic gasoline exposures following Hurricane Sandy. *Prehosp Disaster Med*. 2013;28(6):586-591. doi:10.1017/S1049023X13009023
29. Centers for Disease Control and Prevention. Carbon Monoxide Poisoning: Natural Disasters and Severe Weather. Accessed September 1, 2022. <https://www.cdc.gov/disasters/carbonmonoxide.html>
30. Wang Z, Wu X, Dai W, et al. The Prevalence of Posttraumatic Stress Disorder Among Survivors After a Typhoon or Hurricane: A Systematic Review and Meta-Analysis. *Disaster Med Public Health Prep*. 2019;13(5-6):1065-1073. doi:10.1017/DMP.2019.26
31. Schwartz RM, Gillezeau CN, Liu B, Lieberman-Cribbin W, Taioli E. Longitudinal Impact of Hurricane Sandy Exposure on Mental Health Symptoms. *International Journal of Environmental Research and Public Health* 2017, Vol 14, Page 957. 2017;14(9):957. doi:10.3390/IJERPH14090957
32. Olteanu A, Arnberger R, Grant R, Davis C, Abramson D, Asola J. Persistence of Mental Health Needs among Children Affected by Hurricane Katrina in New Orleans. *Prehosp Disaster Med*. 2011;26(1):3-6. doi:10.1017/S1049023X10000099
33. Schwartz RM, Sison C, Kerath SM, et al. The impact of Hurricane Sandy on the mental health of New York area residents. *Am J Disaster Med*. 2015;10(4):339-346. doi:10.5055/AJDM.2015.0216
34. Lowe D, Ebi KL, Forsberg B. Factors increasing vulnerability to health effects before, during and after floods. *Int J Environ Res Public Health*. 2013;10(12):7015-7067. doi:10.3390/IJERPH10127015
35. Garfin DR, Thompson RR, Holman EA, Wong-Parodi G, Silver RC. Association Between Repeated Exposure to Hurricanes and Mental Health in a Representative Sample of Florida Residents. *JAMA Netw Open*. 2022;5(6):e2217251-e2217251. doi:10.1001/JAMANETWORKOPEN.2022.17251
36. Lowe SR, Rhodes JE, Zwiebach L, Chan CS. The Impact of Pet Loss on the Perceived Social Support and Psychological Distress of Hurricane Survivors. *J Trauma Stress*. 2009;22(3):244. doi:10.1002/JTS.20403
37. Raker EJ, Lowe SR, Arcaya MC, Johnson ST, Rhodes J, Waters MC. Twelve years later: The long-term mental health consequences of Hurricane Katrina. *Soc Sci Med*. 2019;242:112610. doi:10.1016/J.SOCSCIMED.2019.112610
38. Ochi S, Hodgson S, Landeg O, Mayner L, Murray V. Disaster-Driven Evacuation and Medication Loss: a Systematic Literature Review. *PLoS Curr*. 2014;6. doi:10.1371/CURRENTS.DIS.FA417630B566A0C7DFDBF945910EDD96
39. Nogueira LM, Sahar L, Efstathiou JA, Jemal A, Yabroff KR. Association Between Declared Hurricane Disasters and Survival of Patients With Lung Cancer Undergoing Radiation Treatment. *JAMA - Journal of the American Medical Association*. 2019;322(3):269-271. doi:10.1001/jama.2019.7657
40. Stasiewicz AM, Paveglio TB. Preparing for wildfire evacuation and alternatives: Exploring influences on residents' intended evacuation behaviors and mitigations. *International Journal of Disaster Risk Reduction*. 2021;58:102177. doi:10.1016/J.IJDRR.2021.102177